



PATIENT INFORMATION

PERIPHERALLY INSERTED CENTRAL CATHETER (PICC) AMBULATORY PROGRAM ORDER SET

As of January 2022

Contraindications for PICC Insertion

"A peripherally inserted central catheter (PICC) is contraindicated in patients whose veins are compromised (e.g. bruised, infiltrated, phlebotic, sclerosed, corded), those with a history of chronic kidney disease (stage 3 or 4) or patients with an extremity affected by mastectomy, arteriosclerosis, graft, fistula, or radical artery surgery" (Elsevier, 2019).

Allergies: [ ] No Known Allergies OR \_\_\_\_\_

Height (cm): \_\_\_\_\_ Weight (kg): \_\_\_\_\_

CLINICAL INDICATION
[ ] Long-term intravenous therapy
[ ] Chemotherapy/Systemic therapy
[ ] Poor peripheral venous access
[ ] Total Parenteral Nutrition
[ ] Antibiotics (duration or extravasation hazard)
[ ] OTHER:
[ ] Critically ill requiring multi-lumen access
[ ] Frequent need for infusions/blood products
[ ] MAiD (Unable to obtain 2 intravenous saline locks)

LINE REQUESTED
[ ] Single Lumen [ ] Double Lumen

MEDICATIONS
[X] Lidocaine 1% (10mg/mL), 1-2ml intradermally

DIAGNOSTIC IMAGING
[X] Chest x-ray for posterior-anterior (PA), or anterior-posterior (AP) OR Portable
\*For Radiologist to confirm tip placement upon completion of PICC insertion

Table with 3 columns: Date (yyyy/mm/dd), Time, and Provider Name (please print). Includes rows for Provider Signature and Transcriber Name/Signature.