PATIENT INFORMATION



PERIPHERALLY INSERTED CENTRAL CATHETER (PICC) AMBULATORY PROGRAM ORDER SET

As of June 2023

Contraindications for PICC Insertion

"A peripherally inserted central catheter (PICC) is contraindicated in patients whose veins are compromised (e.g. bruised, infiltrated, phlebitic, sclerosed, corded), those with a history of chronic kidney disease (stage 3 or 4) or patients with an extremity affected by mastectomy, arteriosclerosis, graft, fistula, or radical artery surgery" (Elsevier, 2019).

Patient diagnosis:			
Allergies: No Known Allergies OR			
Height (cm): Weight (kg):			
CLINICAL INDICATION			
□ Long-term intravenous therapy □ Critically ill requiring multi-lumen access □ Chemotherapy/Systemic therapy □ Frequent need for infusions/blood products □ Poor peripheral venous access □ MAiD (Unable to obtain 2 intravenous saline locks) *Midline will be used* OTHER:			
LINE REQUESTED			
☐ Single Lumen ☐ Double Lumen ☐ Midline ONLY to be used for clinical indication of MAiD			
MEDICATIONS			
□ Lidocaine 1% (10mg/mL), 1-2ml intradermally			
DIAGNOSTIC IMAGING			
Chest x-ray for posterior-anterior (PA), or anterior-posterior (AP) OR Portable *For Radiologist to confirm tip placement upon completion of PICC insertion			

Date (yyyy/mm/dd):	Time:	Provider Name (please print):
		Provider Signature:
Date (yyyy/mm/dd):	Time:	Transcriber Name (please print):
		Transcriber Signature: